

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> : A01N 37/36, 37/12, 37/04, 37/02 // (A01N 37/36, 57:16, 47:22, 43:84, 43:713, 43:653, 43:54, 37:52) (A01N 37/12, 57:16, 47:22, 43 :84, 43 :713, 43 :653, 43 :54, 37 :22) (A01N 37 /04, 57 :16, 47 :22, 43 :84, 43 :713, 43 :653, 43 :54, 37 :52)</p>	<p>A1</p>	<p>(11) International Publication Number: <b>WO 96/22020</b> (43) International Publication Date: 25 July 1996 (25.07.96)</p>
<p>(21) International Application Number: PCT/GB96/00087 (22) International Filing Date: 17 January 1996 (17.01.96) (30) Priority Data: 9500983.3 19 January 1995 (19.01.95) GB (71) Applicant (for all designated States except US): AGREVO UK LIMITED [GB/GB]; Hauxton, Cambridge CB2 5HU (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): BRIGGS, Geoffrey, Gower [GB/GB]; Chesterford Park, Saffron Walden, Essex CB10 1XL (GB). STOCK, David [GB/GB]; Chesterford Park, Saffron Walden, Essex CB10 1XL (GB). (74) Agent: WELLS, Norman, David; AgrEvo UK Limited, Patent Dept., Chesterford Park, Saffron Walden, Essex CB10 1XL (GB).</p>	<p>(81) Designated States: AL, AM, AU, AZ, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, JP, KG, KR, KZ, LK, LT, LV, MD, MK, MX, NO, NZ, PL, RO, RU, SD, SG, SI, SK, TR, TT, UA, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AZ, BY, KG, KZ, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
<p>(54) Title: PESTICIDAL COMPOSITIONS</p> <p>(57) Abstract</p> <p>The use in a pesticidal composition of at least one aliphatic mono-, di- or triester, optionally substituted by one or more hydroxy groups, said ester(s) having: i) a log octanol-water partition coefficient (log P) of from 5 to 13.1, ii) an equivalent hydrocarbon (EH) value of from 20 to 42; the EH value being calculated as the number of carbon atoms plus five for any hydroxy group present plus 3 for each ester group, and iii) an alcohol moiety comprising at least 2 carbon atoms.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

Pesticidal compositionsField of the invention

This invention concerns pesticidal compositions.

5 Background to the invention

Pesticidal compounds, particularly those used for crop protection, are commonly used in the form of compositions containing one or more co-formulants, for example surfactants. We have found that a new group of  
10 compounds, not previously used in association with pesticidal compounds, can also be used with advantage in association with pesticidal compounds.

Description

In one aspect, the invention provides the use in a  
15 pesticidal composition of at least one aliphatic mono-, di- or triester, optionally substituted by one or more hydroxy groups, said ester(s) having:

- i) a log octanol-water partition coefficient (log P) of from 5 to 13.1,
- 20 ii) an equivalent hydrocarbon (EH) value of from 20 to 42; the EH value being calculated as the number of carbon atoms plus five for any hydroxy group present plus 3 for each ester group, and
- iii) an alcohol moiety comprising at least 2 carbon  
25 atoms.

The term 'ester of the invention' is used herein to indicate an ester having the above properties.

In another aspect the invention provides a  
pesticidal composition which comprises a pesticidal  
30 compound and at least one ester of the invention.

In a further aspect the invention provides a method of combating pests at a locus infested or liable to be infested therewith which comprises applying to said locus an effective amount of a pesticidal composition

which comprises a pesticidal compound and at least one ester of the invention.

5 The log P value of an ester as that term is used herein is that calculated from the structure of the compound using the clog P Program, Pomona College Med Chem release 3.54, January 1988, provided by Daylight Chemical Information Systems Inc, Claremont, California. The values generated by this program correlate well in general with those determined experimentally by methods well known to those skilled in the art, but such methods are difficult to apply accurately to compounds having log P values within the scope of this invention, and we therefore use the calculated values.

10 The EH value can be simply calculated from the structure of each compound. Sometimes, however, the esters of the invention occur or are presented as mixtures of individual compounds, and the mixture will therefore include compounds of different EH values.

15 The ratio of EH value to log P value is preferably in the range 2.4:1 to 4.15:1.

20 Preferred esters of the invention are succinate diesters, lactic acid carboxylic esters, and esters resulting from esterification of a saturated acid of at least 12 carbon atoms with a branched chain alcohol.

25 Examples of suitable esters of the invention include propylene glycol di-ester of coconut fatty acids (C8 ester log P=7.022, EH=25; C10 ester log P=9.138, EH=29), isopropyl palmitate (log P=8.386, EH=22), isopropyl myristate (log P=7.328, EH=20), 2-ethylhexyl 12-hydroxystearate (log P=9.882, EH=34), dioctyl succinate (log P=7.771, EH=26), 2-ethylhexyl palmitate (log P=11.03, EH=27), myristyl lactate (log P=6.556, EH=25), lauryl lactate (log P=5.598, EH=23), and triglycerides of C<sub>8</sub>-C<sub>10</sub>-coconut fatty acids (C8 ester log

P=9.89, EH=36; C10 ester log P=13.064, EH=42).

We have found that the esters of the invention, which have little or no pesticidal activity in their own right, surprisingly aid the penetration of the  
5     pesticidal compound into plant or pest tissue, thereby promoting enhanced activity, particularly of pesticidal compounds whose activity depends on such penetration.

The pesticidal compound is especially one disclosed in The Pesticide Manual, 10th edition, published by Crop  
10     Protection Publications, and may for example be a herbicide, fungicide, insecticide or acaricide.

It is preferably a phytopathogenic fungicide selected from:

- (i)     a conazole steroid demethylation inhibitor;
- 15     (ii)    a steroid reduction inhibitor based on a 1-[3-(4-tert-butylphenyl)-2-methylpropyl] group which is attached via the N-atom to piperidine or 2,6-dimethylmorpholine
- (iii)   a dithiocarbamate fungicide
- 20     (iv)    a phthalimide fungicide in which a chloroalkylthio group is attached via the N-atom to the optionally hydrogenated phthalimide group.
- (v)     an anilide fungicide
- (vi)    an mbc fungicide.
- 25     (vii)   a carbamate fungicide
- (viii)  a copper compound fungicide
- (ix)    a tin compound fungicide
- (x)     a strobilurin type fungicide,
- (xi)    a 2-anilinopyrimidine fungicide or
- 30     (xii)   a fungicide selected from the group consisting of chlorothalonil, dimethomorph, fenpiclonil, fluazinam, hymexazol, nuarimol, pencycuron, pyrifenox, thicyofen, probenazole, pyroquilon, tricyclazole, quaternary ammonium compounds,

fludioxonil, 5-chloro-2-methyl-4-isothiazolin-3-one, 2-methyl-4-isothiazolin-3-one (and mixtures of these two), furmecyclox, 3-iodo-2-propynyl butylcarbamate and sulfur.

5 Conazoles are as defined in ISO standard 257, ie compounds based on imidazole or 1,2,4-triazole and containing a halogenated phenyl group. Examples include prochloraz (and its metal complexes, especially the manganese or copper complex), propiconazole,  
10 flusilazole, hexaconazole, tebuconazole, difenoconazole, bromuconazole, cyproconazole, diniconazole, fenbuconazole, imibenconazole, furconazole, tetraconazole, myclobutanil, penconazole, fluquinconazole, azaconazole, imazalil, triflumizole,  
15 epoxiconazole, triticonazole, metconazole and the fungicide having the code No SSF 109.

Examples of type (ii) fungicides include fenpropimorph and fenpropidin.

20 Examples of type (iii) fungicides include mancozeb and thiram.

Examples of type (iv) fungicides include folpet, captafol and captan.

Examples of type (v) fungicides include  
25 a) 3',5'-dichloroanilide fungicides in which the anilino nitrogen comprises a ring carrying two oxo substituents, in positions adjacent the nitrogen, eg iprodione, vinclozolin or procymidone, or  
b) acetanilide fungicides, eg metalaxyl or ofurace,  
c) sulfanilide fungicides, eg dichlofluanid,  
30 d) benzanilide fungicides, eg flutolanil, and  
e) heteroarylanilide fungicides, eg thifluzamide.

Examples of type (vi) fungicides include carbendazim, benomyl and thiophanate-methyl.

Examples of type (vii) fungicides include

diethofencarb and propamocarb.

Examples of type (viii) fungicides include Bordeaux mixture, oxine-copper, copper oxychloride and copper naphthenate.

5        Examples of type (ix) fungicides include tributyltin oxide and tributyltin naphthenate.

10        Strobilurine type fungicides (type (x) fungicides) are methyl esters of arylacetic acid in which the acetic acid also carries a methoxymethylene or methoxyimino substituent. The aryl group is usually a 2-substituted phenyl group. Examples of such compounds are those disclosed in a wide number of patent applications, including EPs 178826, 203606, 203608, 206523, 229974, 226917, 242070, 242081, 243012, 243014, 251082, 256667, 15        260794, 260832, 267734, 270252, 273572, 274825, 278595, 291196, 299694, 307101, 307103, 310954, 312221, 312243, 329011 and 336211. Specific compounds are those having the code Nos BAS 490F and ICIA 5504.

20        Examples of type (xi) fungicides include pyrimethanil, mepanipyrim and cyprodinil.

Examples of insecticides which may be used include amitraz, triazophos and formetanate.

Examples of acaricides which may be used include clofentezine.

25        The names quoted for these compounds are the non-proprietary common names and the chemical structures can be found for example by reference to The Pesticide Manual.

30        The invention is particularly useful for pesticidal compounds of high melting point and/or low solubility in organic solvents. It is particularly applicable to fluquinconazole.

The esters of the invention may be incorporated in conventional concentrate formulations of the pesticidal

compound (eg suspension concentrates, suspoemulsions or solid formulation types such as water dispersible granules), which are diluted with water prior to application, or they may be added to the pesticidal composition just prior to use.

It may be necessary and/or desirable for the pesticidal composition to include small quantities of solvent and/or surfactant, especially a non-ionic surfactant, and other additives such as fatty acids, to improve the emulsifiability of the esters of the invention. Such modifications are well within the competence of those skilled in the art.

The esters of the invention preferably comprise from 2 to 90% by weight of a concentrate formulation.

The esters of the invention are preferably applied in a diluted formulation at a concentration of 20g to 2000g, preferably 50g to 500g, per 100 litres.

The mixtures can be used for those applications where the pesticide would normally be used. In the case of fluquinconazole, for example, it is preferably used in the direct treatment of growing crops, such as cereals and fruit, or in seed treatments.



The following Examples illustrate the use of esters of the invention, in which:

"Crodamol OHS" is a trade name for 2-ethylhexyl 12-hydroxystearate;

5 "Crodamol PC" is a trade name for propylene glycol diester of coconut fatty acids;

"Crodamol IPP" is a trade name for isopropyl palmitate;

10 "Crodamol IPM" is a trade name for isopropyl myristate;

"Crodamol OHS" is a trade name for 2-ethylhexyl 12-hydroxystearate;

"Crodamol OSU" is a trade name for dioctyl succinate;

15 "Crodamol OP" is a trade name for 2-ethylhexyl palmitate;

"Crodamol ML" is a trade name for myristyl lactate;

"Crodamol LL" is a trade name for lauryl lactate.

Example 1

20 In a glasshouse wheat plants were inoculated with *Erysiphe graminis*. Three days after inoculation the plants were sprayed separately with:

(a) fluquinconazole (FQ), obtained by diluting with water to the desired concentration, a wettable granule  
25 comprising 25% by weight active ingredient,

(b) Crodamol OHS, or

(c) Crodamol PC,

and with a mixture of (a) with (b) or (c) containing about 10% of an emulsifier.

30 8 replicates were used for each treatment. After 4 days, the plants were assessed for the control of sporulation of the disease achieved by the treatments.

The individual components (a), (b) and (c) alone had no effect on sporulation.

Results for the mixtures are given in the following table:

AI	g/ha	(b) (%v/v)	(c) (%v/v)	% control of sporulation
FQ	100	0.4		95.8
FQ	100	0.2		80
FQ	100		0.4	96.9
FQ	100		0.2	89.2

#### Example 2

In a glasshouse wheat plants were inoculated with *Erysiphe graminis*. One day after inoculation the plants were sprayed with:

- (a) an active ingredient (AI), alone and with
- (b) Crodamol OHS, or
- (c) Crodamol PC.

8 replicates were used for each treatment, and each treatment was made by an automatic sprayer delivering the equivalent of 200 l/ha of formulation. After 7 days, the plants were assessed for the control of the disease achieved by the treatments.

Results are given in the following table, in which:

- EC is epoxiconazole
- BC is bromuconazole
- TC is tebuconazole
- FP is fenpropimorph
- PM is pyrimethanil
- CP is cyprodinil.

AI	g/ha	(b) (%v/v)	(c) (%v/v)	% control
EC	31.25	0.4		97.9
EC	31.25	0.2		98.2
EC	31.25		0.4	94.8
EC	31.25		0.2	98.2

9

	EC	31.25			82.6
	BC	62.5	0.4		94.6
	BC	62.5	0.2		89.4
	BC	62.5		0.4	94.8
5	BC	62.5		0.2	96.1
	BC	62.5			77.9
	TC	31.25	0.4		96.4
	TC	31.25	0.2		96.1
	TC	31.25		0.4	94.6
10	TC	31.25		0.2	95.8
	TC	31.25			87.8
	FP	93.75	0.4		62.1
	FP	93.75	0.2		73.0
	FP	93.75		0.4	75.1
15	FP	93.75		0.2	67.8
	FP	93.75			10.1
	PM	800	0.4		54.3
	PM	800	0.2		51.9
	PM	800		0.4	56.8
20	PM	800		0.2	52.2
	PM	800			28.9
	CP	375	0.4		100
	CP	375	0.2		100
	CP	375		0.4	98.4
25	CP	375		0.2	100
	CP	375			87.0
	Untreated Control				0

Example 3

30 Two week old bean plants (Vicia faba) infested with Aphis fabae were sprayed at a volume equivalent to 1000 l/ha with

- (a) an active ingredient (AI), alone and with  
 (b) Crodamol OHS,  
 as set out in the table below.

10

After drying, the plants were placed in a greenhouse, and were assessed 3 days later relative to an untreated control.

Results are given in the following table, in which:

5

AZ is amitraz

TP is triazophos

FM is formetanate.

	AI	g/ha	(b) (%v/v)	% Mortality
	AZ	300	-	0
10	AZ	300	0.3	50
	AZ	100	-	0
	AZ	100	0.3	23
	AZ	30	-	0
	AZ	30	0.3	10
15	TP	300	-	35
	TP	300	0.3	65
	TP	100	-	0
	TP	100	0.3	45
	TP	30	-	0
20	TP	30	0.3	35
	FM	300	-	0
	FM	300	0.3	25

#### Example 4

25 Four week old bean plants (Phaseolus vulgaris) were infested with Tetranychus urticae. After 1 day and after 2 days, the motile stages were blown away, and the plants and remaining eggs were sprayed by hand to runoff with

- 30 (a) clofentezine (CZ) (3g/hl) alone,  
 (b) 0.3% of the esters of the invention specified below, alone, and  
 (c) mixtures of (a) and (b).

After drying, the plants were placed in a greenhouse. 10 days later, the plants were assessed for

11

ovicidal activity relative to an untreated control

Results are given in the following table:

	<u>Ester</u>	<u>% Mortality</u>	<u>% Mortality</u>
		<u>with CZ</u>	<u>without CZ</u>
5	Crodamol IPP	99	0
	Crodamol IPM	90	0
	Crodamol OP	98	8
	Crodamol LL	97	0
	Crodamol PC	63	0
10	Crodamol OHS	80	15
	None	50	0

Example 5

The penetration efficacy of various esters of the invention was demonstrated by applying an acetone/water solution of radio-labelled fluquinconazole (0.05% w/v) both alone and containing the esters specified below to vine plants using a microapplicator. After 5 days, the fluquinconazole remaining on the surface of the plants was recovered and determined quantitatively, the difference between the percentage remaining and 100% being indicative of the amount of compound penetrating the plant tissue. Results are given in the following table:

	<u>Ester</u>	<u>%w/v</u>	<u>Surface %</u>
25	Crodamol PC	0.5	27
	Crodamol IPP	0.5	5
	Crodamol IPM	0.5	10
	Crodamol OHS	0.5	19
	Crodamol OSU	0.5	28
30	Crodamol OP	0.5	26
	Crodamol ML	0.5	10
	Crodamol LL	0.5	22
	None	-	>90

CLAIMS

1. The use in a pesticidal composition of at least one aliphatic mono-, di- or triester, optionally substituted by one or more hydroxy groups, said ester(s) having:
  - i) a log octanol-water partition coefficient (log P) of from 5 to 13.1,
  - ii) an equivalent hydrocarbon (EH) value of from 20 to 42; the EH value being calculated as the number of carbon atoms plus five for any hydroxy group present plus 3 for each ester group, and
  - iii) an alcohol moiety comprising at least 2 carbon atoms.
2. The use according to claim 1 of at least one ester having a ratio of EH value to log P value in the range 2.4:1 to 4.15:1.
3. The use according to claim 1 or claim 2 of at least one ester which is a succinate diester, a lactic acid carboxylic ester, or an ester resulting from esterification of a saturated acid of at least 12 carbon atoms with a branched chain alcohol.
4. The use according to any of claims 1 to 3 of at least one ester selected from propylene glycol di-ester of coconut fatty acids, isopropyl palmitate, isopropyl myristate, 2-ethylhexyl 12-hydroxy-stearate, dioctyl succinate, 2-ethylhexyl palmitate, myristyl lactate, lauryl lactate, and triglycerides of C<sub>8</sub>-C<sub>10</sub>-coconut fatty acids.
5. The use according to any of claims 1 to 4 in which the pesticidal composition is a phytopathogenic fungicidal composition.

6. The use according to claim 5 in which the composition comprises fluquinconazole as active ingredient.
- 5 7. A pesticidal composition which comprises a pesticidal compound and at least one ester as defined in any of claims 1 to 4.
8. A composition according to claim 7 which comprises at least one ester as defined in any of claims 1 to 4 in a concentration of from 2 to 90% by weight.
- 10 9. A method of combating pests at a locus infested or liable to be infested therewith which comprises applying to said locus an effective amount of a pesticidal composition which comprises a pesticidal compound and at least one ester of the invention as defined in any of claims 1 to 4.
- 15 10. A method according to claim 9 in which the ester of the invention is applied in a concentration of from 20g to 2000g per 100 litres of the applied composition.

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 96/00087

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A01N37/36 A01N37/12 A01N37/04 A01N37/02 //(A01N37/36,  
57:16,47:22,43:84,43:713,43:653,43:54,37:52), (A01N37/12,57:16,  
47:22,43:84,43:713,43:653,43:54,37:22), (A01N37/04,57:16,47:22,

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE WPI Week 7528 Derwent Publications Ltd., London, GB; AN 75-46209W [28] XP002003965 & DD,A,112 891 (HOECHST) , 12 May 1975 see abstract	1-5,7-10
X	DE,A,23 03 757 (HOECHST AG) 15 August 1974 see page 2, paragraph 2 see page 3, paragraph 2 see page 3, paragraph 4 see page 4, paragraph 4	1-5,7-10
X	DE,A,42 39 181 (HOECHST AG) 5 May 1994 see page 2, line 17 - line 24 see page 3, line 1 see page 3, line 9 - line 10	1-4,7-10
-/-		

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

\*A\* document member of the same patent family

Date of the actual completion of the international search

28 May 1996

Date of mailing of the international search report

11.06.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+ 31-70) 340-3016

Authorized officer

Lamers, W



## INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 96/00087

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 43:84,43:713,43:653,43:54,37:52)

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	PATENT ABSTRACTS OF JAPAN vol. 5, no. 160 (C-075), 15 October 1981 & JP,A,56 092207 (NISSHIN OIL MILLS), 25 July 1981, see abstract	1-4,7-10
X	DE,A,32 47 050 (BAYER AG) 20 June 1984 see page 11, paragraph 4 - page 12, paragraph 1 see page 18, line 12 - line 13 see page 18, line 15 - page 19, line 11	1-4,7-10
Y	see page 11, paragraph 2	4
Y	FR,A,2 347 052 (BAYER AG) 4 November 1977 see page 3, line 6 - page 4, line 21 see page 5, line 33 - page 6, line 16	4
	--- -/--	

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- \*Z\* document member of the same patent family

Date of the actual completion of the international search

Date of mailing of the international search report

11.06.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+ 31-70) 340-3016

Authorized officer

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 96/00087

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP,A,0 295 411. (SANSHO CO ;EISAI CO LTD (JP)) 21 December 1988 see page 2, line 30 - line 34 see page 2, line 53 - page 3, line 1 ---	4
Y	DATABASE WPI Week 9351 Derwent Publications Ltd., London, GB; AN 93-408837 [51] XP002003457 & JP,A,05 306 223 (MAEDA) , 19 November 1993 see abstract ---	4
X	EP,A,0 579 052 (SCHOENHERR JOERG PROF DR) 19 January 1994 see page 2, line 1 - line 4 see page 2, line 41 - line 46 see page 3, line 26 - line 29 see page 3, line 46 - line 53 see page 4, line 33 - line 38 ---	1,2,5, 7-10
X	DATABASE WPI Week 9316 Derwent Publications Ltd., London, GB; AN 93-129070 [16] XP002003458 & JP,A,05 065 201 (DOWELANCO) , 19 March 1993 see abstract ---	1-4,7-10
X	EP,A,0 045 424 (BAYER AG) 10 February 1982 see page 2, paragraph 4 see page 4, paragraph 4 see page 5, paragraph 1 - paragraph 2 ---	1-4,7-10
X	EP,A,0 128 351 (BAYER AG) 19 December 1984 see page 1, line 1 - line 6 see page 2, line 9 - page 3, line 20 ---	1-4,7-10
A	CHEMICAL ABSTRACTS, vol. 118, no. 19, 10 May 1993 Columbus, Ohio, US; abstract no. 185737, P.E.RUSSELL ET AL.: "Fluquinconazole, a novel broad-spectrum fungicide for foliar application" XP002003455 see abstract & BRIGHTON CROP PROT.CONF.--PESTS DIS., no. 1, 1992, pages 411-418, --- -/--	6

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 95/00087

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>CHEMICAL ABSTRACTS, vol. 105, no. 21, 24 November 1986 Columbus, Ohio, US; abstract no. 185721, A.CHAMEL: "Foliar absorption of herbicides: study of the cuticular penetration using isolated cuticles" XP002003456 see abstract &amp; PHYSIOL.VEG., vol. 24, no. 4, 1986, pages 491-508, -----</p>	1

# INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/GB 96/00087

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0045424	10-02-82	DE-A- 3029426	11-03-82
		AR-A- 228459	15-03-83
		AU-B- 540763	06-12-84
		AU-B- 7364081	11-02-82
		JP-B- 1031481	26-06-89
		JP-C- 1688883	11-08-92
		JP-A- 57054101	31-03-82
		US-A- 5059593	22-10-91
EP-A-0128351	19-12-84	DE-A- 3317823	22-11-84
		AU-B- 2806384	22-11-84
		JP-A- 59212403	01-12-84

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 96/00087

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE-A-2303757	15-08-74	BE-A- 818086	27-01-75
DE-A-4239181	05-05-94	CN-A- 1088049	22-06-94
		EP-A- 0596316	11-05-94
		JP-A- 6219901	09-08-94
		ZA-A- 9308166	06-06-94
DE-A-3247050	20-06-84	AU-B- 2237983	28-06-84
		EP-A, B 0113857	25-07-84
		JP-A- 59118701	09-07-84
		US-A- 4626274	02-12-86
FR-A-2347052	04-11-77	DE-A- 2614841	20-10-77
		AT-B- 354182	27-12-79
		AU-B- 506372	20-12-79
		AU-B- 2386177	12-10-78
		BE-A- 853303	06-10-77
		CA-A- 1106284	04-08-81
		CH-A- 639822	15-12-83
		GB-A- 1527584	04-10-78
		JP-C- 1451072	25-07-88
		JP-A- 52122619	15-10-77
		JP-B- 62051932	02-11-87
		NL-A- 7703729	10-10-77
		SE-B- 428173	13-06-83
		SE-A- 7703975	07-10-77
		US-A- 4096262	20-06-78
EP-A-0295411	21-12-88	JP-A- 1052716	28-02-89
		JP-B- 7020866	08-03-95
		CA-A- 1310270	17-11-92
		DK-B- 169016	01-08-94
		FI-B- 91361	15-03-94
		NO-B- 175515	18-07-94
		US-A- 5059427	22-10-91
		US-A- 5073375	17-12-91
EP-A-0579052	19-01-94	DE-A- 4319263	05-01-94
		CA-A- 2099631	04-01-94